

Quantum effects in QED_{3+1} with an extra compactified dimension

C. Ccapa Ttira^a, C. D. Fosco^a,
A. P. C. Malbouisson^b
and I. Roditi^b

^a*Centro Atómico Bariloche and Instituto Balseiro
Comisión Nacional de Energía Atómica
8400 Bariloche, Argentina.*

^b*Centro Brasileiro de Pesquisas Físicas - CBPF/MCT
Rua Dr. Xavier Sigaud, 150, 22290-180 Rio de Janeiro, RJ, Brazil*

December 27, 2008

Abstract

We evaluate one-loop effects for QED_{3+1} in the presence of an extra compactified dimension, in a non-trivial vacuum for the gauge field, such that a non-vanishing magnetic flux is encircled along the extra dimension. We obtain the vacuum polarization tensor and evaluate the exact parity breaking term.

Keywords: *Field theory in extra dimensions, Compactification.*

PACS: 11.10.Kk, 12.20.Ds

Quantum field theory models with compactified dimensions have been proposed to investigate many different physical phenomena. These range from finite size effects in critical phenomena [1], to particle physics models where, since the pioneering works of Kaluza and Klein [2], extra dimensions have played an important role to search for methods to unify the fundamental interactions and to describe a wide range of physical situations; to name a few: alternative solutions to the hierarchy problem [3, 4], physics at the TeV scale [5], and particle physics models where the extra dimensions play an important role to implement the unification of fundamental interactions (see, for example, [6, 7, 8] and references therein) .

An interesting yet simple example of this, corresponds to the compactification of one dimension in an $\mathbb{R}^D$ Euclidean spacetime, such that the topology of the resulting manifold $\mathcal{M}$ is that of $\mathbb{R}^{D-1} \times \mathcal{S}_1$, i.e., ‘circular compactification’. Although the interesting features that emerge in this situation have been studied using several different techniques in the literature, one can take advantage of a (formal) common feature; indeed, they share many properties with the imaginary-time formulation of quantum field theory at finite temperature [9, 10]. This allows one, for example, to take advantage of the many well-known methods and results developed in this context, such as Feynman diagrams and renormalization techniques, to import them to the case under consideration.

From a topological point of view, the imaginary-time formalism in a D -dimensional space-time at finite temperature is equivalent to a path-integral evaluated on the manifold $\mathbb{R}^{D-1} \times \mathcal{S}_1$, where $\mathcal{S}_1$ is a circle of circumference $\beta = 1/T$. Since in Euclidean path integral (imaginary) time and spatial coordinates are on the same footing, the Matsubara prescription can be thought of as a mechanism to deal with spatial compactifications where the compactification length L plays the role of β [11]. This has been developed by considering both simply or non simply-connected D -dimensional manifolds with a topology $\mathbb{R}_d^D = \mathbb{R}^{D-d} \times \mathcal{S}_{l_1} \times \mathcal{S}_{l_2} \cdots \times \mathcal{S}_{l_d}$, with l_1 corresponding to the inverse temperature and $l_2, \dots, l_d$ to the compactification of $d - 1$ spatial dimensions¹.

For just one compactified dimension (imaginary time or a spatial dimension) the Feynman rules are modified, the most characteristic new feature is the Matsubara prescription for momentum integrals,

$$\int \frac{dk_s}{2\pi} \rightarrow \frac{1}{\xi} \sum_{n=-\infty}^{+\infty} ; \quad k_i \rightarrow \frac{2n\pi}{\xi} , \quad (1)$$

where k_s amounts to the momentum component corresponding to the compactified dimension, while ξ equals β or L , for the finite temperature and compactified spatial dimension cases, respectively.

Some of these ideas have recently attracted renewed interest, as a way to investigate the electroweak transition and baryogenesis. For instance, a recent investigation of the electroweak phase transition has been presented in [13], in the context of a $4 + 1$ -dimensional theory with a compactified dimension. This model involves charged scalar bosons coupled to a gauge field, and makes use of the Hosotani mechanism [14]. Taking the order parameter to be proportional to the non-vanishing component of the gauge field along

¹This case has been considered, within the context of spontaneous symmetry breaking, in [12].

the compactified coordinate, the authors find a first-order transition with a strength inversely proportional to the Higgs mass.

Within the previous general framework, we here take a slightly different route, to investigate one-loop effects for QED_{3+1} with an extra compactified dimension, in a non-trivial vacuum for the gauge field, with the non-vanishing component of the gauge field lying also along the extra dimension.

The system we shall deal with may be conveniently defined in terms of an Euclidean action, S , which has the structure:

$$S(\mathcal{A}; \bar{\Psi}, \Psi) = S_g(\mathcal{A}) + S_f(\mathcal{A}; \bar{\Psi}, \Psi), \quad (2)$$

where S_g and S_f denote the $U(1)$ gauge field and fermionic actions, respectively. The former is assumed to have a standard Maxwell form, namely:

$$S_g(\mathcal{A}) = \frac{1}{4} \int d^5x \mathcal{F}_{\alpha\beta} \mathcal{F}_{\alpha\beta}, \quad (3)$$

with $\mathcal{F}_{\alpha\beta} \equiv \partial_\alpha \mathcal{A}_\beta - \partial_\beta \mathcal{A}_\alpha$, where we adopted the convention that indices from the beginning of the Greek alphabet ($\alpha, \beta, \dots$) label all the coordinates of the spacetime manifold, and therefore run from 0 to 4. Since we will be specially interested in the model as it is seen from a $3+1$ dimensional point of view, we shall also use another convention: indices from the middle of the Greek alphabet ($\mu, \nu, \dots$) are reserved for the $3+1$ -dimensional spacetime coordinates while, when this notation is used, the extra dimension coordinate shall be denoted by s . Then:

$$\alpha = 0, 1, 2, 3, 4, \quad \mu = 0, 1, 2, 3, \quad d^5x \equiv d^{3+1}x dx_4 = d^{3+1}x ds, \quad (4)$$

and x will be assumed to denote the $3+1$ coordinates x_μ , unless explicit indication on the contrary. The extra dimension is assumed to be compactified with a radius R , so that $s \sim s + L$, $L = 2\pi R$.

On the other hand, the Dirac action, S_f , is given by

$$S_f(\bar{\Psi}, \Psi; \mathcal{A}) = \int d^{3+1}x ds \bar{\Psi}(x, s) (\mathcal{D} + m) \Psi(x, s) \quad (5)$$

where $\mathcal{D}$ is the $4+1$ dimensional Dirac operator, $\mathcal{D} = \gamma_\alpha D_\alpha$. The covariant derivative $D_\alpha \equiv \partial_\alpha + ig\mathcal{A}_\alpha$ includes a coupling constant g with the dimensions of $(\text{mass})^{-\frac{1}{2}}$. For Dirac's γ -matrices, we assume that $\gamma_s \equiv \gamma_5$, where the latter is the γ_5 matrix for the $3+1$ world.

To proceed, we discuss now the mode expansion and its relation to gauge invariance. To that end, we follow [15], where this issue is discussed at length, albeit in the finite temperature theory context, in the Matsubara formulation

of thermal field theory. Due to the formal analogy with this situation, a quite straightforward procedure allows us to adapt the results derived there to our case. The necessary changes that follow from the fact that our compactified dimension is spatial rather than temporal are taken into account by using (1). In that analogy, the length L plays the same role of the inverse temperature in [15]: $L \sim \beta$, $\beta = T^{-1}$.

What follows is a brief review of some of those properties (the ones which are relevant to our study), adapted to our case and conventions. To begin with, the gauge field configuration $\mathcal{A}_\alpha(x, s)$ may be decomposed into its zero (A_α) and non-zero (Q_α) mode components:

$$\mathcal{A}_\alpha(x, s) = L^{-\frac{1}{2}} A_\alpha(x) + Q_\alpha(x, s) , \quad (6)$$

where the two terms in this decomposition may be defined by:

$$A_\alpha(x) = L^{-\frac{1}{2}} \int_0^L ds \mathcal{A}_\alpha(x, s) , \quad (7)$$

and

$$Q_\alpha(x, s) = \mathcal{A}_\alpha(x, s) - L^{-\frac{1}{2}} A_\alpha(x) , \quad (8)$$

so that $\int_0^L ds Q_\alpha(x, s) = 0$. An $L^{-\frac{1}{2}}$ factor has been included in the zero mode term in order to make this field have the usual mass dimensions in 3+1 spacetime dimensions; this property will become useful after dimensional reduction.

The decomposition above finds a natural interpretation when one considers the Fourier expansion of the gauge field along the extra dimension:

$$\mathcal{A}_\alpha(x, s) = L^{-\frac{1}{2}} \sum_{n=-\infty}^{\infty} e^{i\omega_n s} \tilde{\mathcal{A}}_\alpha(x, n) , \quad (9)$$

with $\omega_n \equiv \frac{2\pi n}{L}$, where one identifies:

$$A_\alpha(x) = \tilde{\mathcal{A}}_\alpha(x, 0) , \quad Q_\alpha(x, s) = L^{-\frac{1}{2}} \sum_{n \neq 0} e^{i\omega_n s} \tilde{\mathcal{A}}_\alpha(x, n) . \quad (10)$$

Then we dimensionally reduce the theory, what, for the gauge field action, amounts to keeping just the zero mode component of the gauge field. Thus:

$$S_g(\mathcal{A}) \rightarrow S_g(A) = S_g(A_\mu, A_s) , \quad (11)$$

where:

$$S_g(A_\mu, A_s) = \int d^{3+1}x \left[\frac{1}{2} \partial_\mu A_s \partial_\mu A_s + \frac{1}{4} F_{\mu\nu}(A) F_{\mu\nu}(A) \right] , \quad (12)$$

with $F_{\mu\nu}(A) \equiv \partial_\nu A_\mu - \partial_\mu A_\nu$.

Regarding the fermionic action S_f , the reduction amounts to:

$$S_f(\mathcal{A}; \bar{\Psi}, \Psi) \rightarrow S_f(A_\mu, A_s; \bar{\Psi}, \Psi) . \quad (13)$$

The fermionic field is not dimensionally reduced by the simple reason that, in the calculation of the effective gauge field action, its only contribution comes from the fermion loop. That loop may be represented as a series of $3 + 1$ loops, each one with a different mass. Although the contributions of heavier modes may be relatively suppressed, the very fact that there is an infinite number of them forbids us to truncate that series (even if there were a zero mode).

Thus, the following explicit expression for the fermionic action shall be used after dimensional reduction:

$$S_f = \int d^{3+1}x \int_0^L ds \bar{\Psi}(x, s) (\not{D} + \gamma_s D_s + m) \Psi(x, s) \quad (14)$$

where

$$\not{D} = \gamma_\mu (\partial_\mu + ieA_\mu) \quad D_s = \partial_s + ieA_s . \quad (15)$$

We have introduced a new, dimensionless coupling constant $e \equiv gL^{-\frac{1}{2}}$, which shall play the role of the electric charge in $3 + 1$ dimensions.

As explained in [15], when considering the form of the gauge transformations in terms of the decomposition into zero and non-zero modes, one finds that A_μ transforms as a standard gauge field (in $3 + 1$ dimensions):

$$\delta A_\mu(x) = \partial_\mu \alpha(x) \quad (16)$$

while its extra dimensional component A_s , a scalar from the $3+1$ -dimensional point of view, is shifted by a constant:

$$\delta A_s(x) = \Omega . \quad (17)$$

The constant Ω has to be of the form $\Omega = \frac{2\pi k}{Le}$, where k is an integer, since the gauge field is coupled to a (charged) fermionic field, whose transformation law under simultaneous action of the previous gauge transformations is:

$$\begin{aligned} \Psi(x, s) &\rightarrow e^{-ie[\alpha(x) + \Omega s]} \Psi(x, s) \\ \bar{\Psi}(x, s) &\rightarrow e^{ie[\alpha(x) + \Omega s]} \bar{\Psi}(x, s) . \end{aligned} \quad (18)$$

We now define the part of the effective action that only depends on the (dimensionally reduced) gauge field, $\Gamma(A)$,

$$\Gamma(A) \equiv \Gamma(A; \bar{\Psi}, \Psi) \Big|_{\bar{\Psi}=\Psi=0} , \quad (19)$$

where $\Gamma(A; \bar{\Psi}, \Psi)$ is the full effective action. The functional $\Gamma(A)$ allows one to derive *1PI* functions containing only A_μ , A_s external lines. The former have an immediate $3 + 1$ dimensional interpretation, while the latter shall be assumed to have a constant (but otherwise arbitrary) value, which is determined by a condition which is external to the model.

On the other hand, at the one-loop order, the only non-trivial term comes from the fermionic loop:

$$\Gamma(A) = \Gamma^{(0)}(A) + \Gamma^{(1)}(A) + \dots \quad (20)$$

where $\Gamma^{(0)}(A) = S_g(A)$ and

$$e^{-\Gamma^{(1)}(A)} = \int \mathcal{D}\Psi \mathcal{D}\bar{\Psi} e^{-S_f(A; \bar{\Psi}, \Psi)} . \quad (21)$$

We shall focus on the effective action for the gauge field components A_μ that have a direct physical interpretation from a $3 + 1$ -dimensional perspective. Regarding the scalar component, A_s , as we have said above, it will be assumed to yield a non-vanishing flux:

$$e \int_0^L ds A_s = \phi \quad (22)$$

where ϕ is a constant. This condition may be conveniently solved by means of a constant A_s :

$$A_s = \frac{\phi}{eL} , \quad (23)$$

which is the gauge fixing that we shall assume. Note that, since the gauge transformations shift A_s by an integer multiple of $\frac{2\pi}{eL}$, we may fix the value of ϕ to the fundamental region:

$$0 \leq \phi < 2\pi , \quad (24)$$

which we shall assume in what follows.

We then proceed to Fourier expand the fermionic fields along the s coordinate:

$$\begin{aligned} \Psi(x, s) &= L^{-\frac{1}{2}} \sum_{n=-\infty}^{\infty} e^{i\omega_n s} \psi_n(x) \\ \bar{\Psi}(x, s) &= L^{-\frac{1}{2}} \sum_{n=-\infty}^{\infty} e^{-i\omega_n s} \bar{\psi}_n(x) , \end{aligned} \quad (25)$$

and insert this into the functional expression for $\Gamma^{(1)}(A)$, to obtain:

$$S_f = \sum_{n=-\infty}^{n=+\infty} \int d^{3+1}x \bar{\psi}_n(x) \left(\not{D} + i\gamma_s(\omega_n + \frac{\phi}{L}) + m \right) \psi_n(x). \quad (26)$$

Under the same expansion, the fermionic measure factorizes:

$$\mathcal{D}\Psi \mathcal{D}\bar{\Psi} = \prod_{n=-\infty}^{n=+\infty} \mathcal{D}\psi_n(x) \mathcal{D}\bar{\psi}_n(x), \quad (27)$$

and, finally, the Euclidean action corresponding to each mode n may be equivalently written as follows

$$\begin{aligned} & \int d^{3+1}x \bar{\psi}_n(x) \left(\not{D} + i\gamma_s(\omega_n + \frac{\phi}{L}) + m \right) \psi_n(x) \\ &= \int d^{3+1}x \bar{\psi}_n(x) (\not{D} + M_n e^{-i\varphi_n \gamma_5}) \psi_n(x) \end{aligned} \quad (28)$$

with

$$M_n \equiv \sqrt{m^2 + (\omega_n + \phi/L)^2}, \quad \varphi_n = \arctan\left(\frac{\omega_n + \phi/L}{m}\right). \quad (29)$$

The existence of a γ_5 term means that parity symmetry will generally be broken; to study that phenomenon more clearly, we perform a change in the fermionic variables that gets rid of the dependence in γ_5 ,

$$\psi_n(x) \rightarrow e^{-i\gamma_5 \phi_n/2} \psi_n(x), \quad \bar{\psi}_n(x) \rightarrow \bar{\psi}_n(x) e^{-i\gamma_5 \phi_n/2}, \quad (30)$$

after which the mode labelled by n has the action:

$$\int d^{3+1}x \bar{\psi}_n(x) (\not{D} + M_n e^{-i\varphi_n \gamma_5}) \psi_n(x). \quad (31)$$

This chiral rotation in the $3+1$ Euclidean fermionic variables induces, however an anomalous Jacobian $\mathcal{J}_n$ for each mode. Then, $\Gamma^{(1)}$ may be written as follows:

$$e^{-\Gamma^{(1)}(A)} = \prod_{n=-\infty}^{+\infty} \left[\mathcal{J}_n e^{-\Gamma_{3+1}^{(1)}(A, M_n)} \right], \quad (32)$$

where

$$\mathcal{J}_n = \exp\left(\frac{ie^2}{16\pi^2} \phi_n \int d^{3+1}x \tilde{F}_{\mu\nu} F_{\mu\nu}\right), \quad (33)$$

with $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu\rho\lambda}F_{\rho\lambda}$, and $\Gamma_{3+1}^{(1)}(A, M_n)$ is the one-loop fermionic contribution to the effective action, for a fermion whose mass is M_n , in $3 + 1$ dimensions. Of course, it may be expressed as a fermionic determinant:

$$e^{-\Gamma_{3+1}^{(1)}(A, M_n)} = \det(\not{D} + M_n) . \quad (34)$$

Then, we arrive to a general expression for the one loop effective action,

$$\Gamma^{(1)}(A) = \Gamma_e^{(1)}(A) + \Gamma_o^{(1)}(A) \quad (35)$$

where the e and o subscripts stand for the even and odd components (regarding parity transformations) and are given by

$$\Gamma_e^{(1)}(A) = \sum_{n=-\infty}^{\infty} \Gamma_{3+1}^{(1)}(A, M_n) \quad (36)$$

and

$$\Gamma_o^{(1)}(A) = - \sum_{n=-\infty}^{\infty} \ln \mathcal{J}_n , \quad (37)$$

respectively.

The parity conserving part of the effective action may be obtained by performing the sum of the required QED_{3+1} object, with an n -dependent mass, M_n . We shall focus on that part of $\Gamma_e^{(1)}$ that contributes to the vacuum polarization tensor for the A_μ gauge field components. Since we are not interested in response functions which involve the s component of the currents, it is useful to define:

$$\Gamma_e^{(1)}(A_\mu) \equiv \Gamma_e^{(1)}(A_\mu, A_s) - \Gamma_e^{(1)}(0, A_s) . \quad (38)$$

Note that $\Gamma_e^{(1)}(0, A_s) \equiv \Gamma_s(A_s)$ does not contribute to response functions involving A_μ , although it can be used to study the fermion-loop corrections to an A_s effective potential. The explicit form of this function is [15]:

$$\Gamma_s(A_s) = -2L \int d^{3+1}x \int \frac{d^4k}{(2\pi)^4} \ln [\cosh(Lk) + \cos\phi] . \quad (39)$$

The vacuum polarization tensor $\Pi_{\mu\nu}$ is obtained from the quadratic term in a functional expansion in the gauge field:

$$\Gamma_e^{(1)}(A_\mu) = \frac{1}{2} \int d^{3+1}x \int d^{3+1}y A_\mu(x) \Pi_{\mu\nu}(x, y) A_\nu(y) + \dots \quad (40)$$

It is then sufficient to resort to the analogous expansion for the $3 + 1$ dimensional effective action,

$$\Gamma_{3+1}^{(1)}(A, M_n) = \frac{1}{2} \int d^{3+1}x \int d^{3+1}y A_\mu(x) \Pi_{\mu\nu}^{(n)}(x, y) A_\nu(y) + \dots \quad (41)$$

(which is even) so that the vacuum polarization receives contributions from all the modes:

$$\Pi_{\mu\nu}^e = \sum_n \Pi_{\mu\nu}^{(n)}, \quad (42)$$

where $\Pi_{\mu\nu}^{(n)} = \Pi^{(n)}(k^2) \delta_{\mu\nu}^T(k)$, with:

$$\Pi^{(n)}(k^2) = \frac{2e^2}{\pi} \int_0^1 d\beta \beta(1-\beta) \ln \left[1 + \beta(1-\beta) \frac{k^2}{M_n^2} \right], \quad (43)$$

the renormalized scalar part of the vacuum polarization tensor, and the transverse projector is defined by $\delta_{\mu\nu}^T(k) \equiv \delta_{\mu\nu} - k_\mu k_\nu / k^2$.

The sum in (42) may be evaluated using standard finite temperature techniques; this yields the result $\Pi_{\mu\nu}^e = \Pi_e \delta_{\mu\nu}^\perp$, with:

$$\begin{aligned} \Pi_e(k^2) &= \frac{2e^2}{\pi} \int_0^1 d\beta \beta(1-\beta) \\ &\times \ln \left\{ \cos(\phi) - \cosh \left[mL \sqrt{1 + \beta(1-\beta) \frac{k^2}{m^2}} \right] \right\}. \end{aligned} \quad (44)$$

We see that, as a consequence of the sum over modes, the expression above does not satisfy the renormalization condition that follows by imposing the validity of Coulomb's law at long distances. However, it is quite straightforward to impose it now, since we only need to perform a finite renormalization, to obtain a properly renormalized function Π_R , whose explicit form is:

$$\begin{aligned} \Pi_R^e(k^2) &= \Pi_e(k^2) - \Pi_e(0) \\ &= \frac{2e^2}{\pi} \int_0^1 d\beta \beta(1-\beta) \ln[1 + F(k^2)], \end{aligned} \quad (45)$$

with

$$F(k^2) = \frac{\cosh \left[mL \sqrt{1 + \beta(1-\beta) \frac{k^2}{m^2}} \right] - \cosh(mL)}{\cosh(mL) - \cos(\phi)}. \quad (46)$$

It is interesting to note that, even though the theory is 5 dimensional, the vacuum polarization tensor requires, to be renormalized, just one renormalization condition, as in 4 dimensions. Indeed, the superficial degree of divergence, $\delta(\gamma)$, for an $1PI$ Feynman graph γ in QED_5 is

$$\delta(\gamma) = 5 - \frac{3}{2}E_G - 2E_F + \frac{1}{2}V \quad (47)$$

where E_G and E_F are the number of external gauge and fermion lines, respectively, and V is the number of vertices. For the one-loop vacuum polarization tensor, we then have $\delta(\gamma) = 3$, which, taking into account gauge invariance is reduced to 1. Moreover, since the divergent terms can only be even polynomials in the momentum, we are left with a zero degree divergence, i.e., a constant.

Let us now study some immediate properties and consequences that follow from expressions (45) and (46) above. The natural approach is perhaps to look at its predictions for different momentum regimes. Let us thus begin by considering the low momentum regime, namely, $k^2 \ll m^2$. The leading term, $k^2/m^2 \rightarrow 0$ has already been considered, to impose the renormalization condition $\Pi_R^e \rightarrow 0$, which is not actually a prediction, but rather the fact that the model contains Coulomb's law at long distances.

The next-to-leading term already contains a non trivial effect. Indeed, a simple effect that is sensible to the presence of the flux is the strength of the Lamb shift, which is determined by the Darwin term. This can be seen by expanding the renormalized tensor to $(\frac{k}{m})^2$ order in a momentum expansion:

$$\Pi_R(k^2) \sim -\frac{e^2}{30\pi} \left[\frac{mL \sinh(mL)}{\cos(\phi) - \cosh(mL)} \right] \frac{k^2}{m^2}, \quad k^2 \sim 0, \quad (48)$$

which for the Hydrogen atom produces a corrected potential energy:

$$V_{eff}(r) = -\frac{e^2}{4\pi r} - \frac{e^4}{120\pi^2 m^2} \left[\frac{mL \sinh(mL)}{\cosh(mL) - \cos \phi} \right] \delta^{(3)}(\mathbf{r}). \quad (49)$$

The usual correction is obtained when $\phi \rightarrow 0$ and $mL \rightarrow 0$:

$$V_{eff}(r) \rightarrow -\frac{e^2}{4\pi r} - \frac{e^4}{60\pi^2 m^2} \delta^{(3)}(\mathbf{r}). \quad (50)$$

It is interesting to study the shape of the ratio between the corrected and usual strengths of the respective Darwin terms:

$$\xi(mL, \phi) \equiv \frac{2mL \sinh(mL)}{\cosh(mL) - \cos \phi}. \quad (51)$$

The case of a vanishing flux yields simply $\xi(mL, 0) = \frac{\frac{mL}{2}}{\tanh(\frac{mL}{2})}$, which for small values of mL approaches 1, and grows linearly with mL when $mL \gg 1$.

The opposite regime, when the effect of the flux is maximum, corresponds to $\phi = \frac{\pi}{2}$:

$$\xi(mL, \frac{\pi}{2}) = 2mL \tanh(mL). \quad (52)$$

The behaviour in this case is quite different; it tends to zero quadratically for small mL , and also grows linearly in the opposite case, albeit with a different slope.

Let us now consider the would be large-momentum region for the vacuum polarization. This regime will be defined by the condition that $k^2 \gg m^2$, although k (and m), will be assumed to be much smaller than L^{-1} . The latter is enforced in order to say that the mass of the Kaluza-Klein modes is much larger than the photon momentum. Under this assumption, one gets the expression:

$$\Pi_R^e(k^2) \sim \frac{2e^2}{\pi} \int_0^1 d\beta \beta(1-\beta) \ln \left[1 + \beta(1-\beta) \frac{k^2}{m_{eff}^2} \right], \quad (53)$$

where

$$m_{eff} \equiv \frac{2|\sin \frac{\theta}{2}|}{L}. \quad (54)$$

We conclude that, as a consequence of the existence of the non-vanishing flux, the large-momentum behaviour differs from the one that one has in standard *QED*, by the emergence of an effective mass m_{eff} . This mass should, in order not to spoil the known anti-screening effect at short distances, be very small. Since L is assumed to be very small, that can only be achieved with an extremely small θ , namely $\theta \ll 1$. Hence,

$$m_{eff} \equiv \frac{2|\sin \frac{\theta}{2}|}{L} \sim \frac{|\theta|}{L} \ll 1. \quad (55)$$

In natural units, if $L^{-1} \equiv \Lambda$ is the large momentum scale set by the Kaluza-Klein modes, and we want m_{eff} to be much smaller than the electron mass, since only in that situation we recover the expected behaviour for the effective charge at small distances. Then we should have:

$$|\theta| \ll \frac{m}{\Lambda}. \quad (56)$$

Finally, the parity breaking term, Γ_o is simply obtained by taking into account (37) and (33):

$$\Gamma_o = -\frac{ie^2}{16\pi^2} \Phi \int d^{3+1}x \tilde{F}_{\mu\nu} F_{\mu\nu}, \quad (57)$$

where we introduced the factor:

$$\Phi = \sum_{n=-\infty}^{\infty} \phi_n; \quad (58)$$

the sum of this series is well-known [16], the result being:

$$\Phi = \arctan \left[\tanh\left(\frac{mL}{2}\right) \tan(\phi/2) \right] . \quad (59)$$

The possible effects due to this term are more difficult to elucidate, since they would require the existence of non-trivial Abelian gauge field background to manifest themselves. Within the present model, there is no room to accommodate them, except if singular configurations were

To conclude, we enumerate the main points we wished to convey in this letter: by studying the vacuum polarization function in this *QED* model with an extra dimension and flux, physical effects due to the compactification can be found at the level of the vacuum polarization tensor. Indeed, the strongest effect is due to the non-vanishing flux, parametrized by θ , in the large momentum behaviour of the effective charge. We see that θ should be much smaller than the ratio between the electron mass and the (momentum) scale induced by the inverse of the compactification radius in order for this effect to be hidden. Besides, the effect of the non-vanishing flux is maximum when it reaches π . This is to be expected, since in that case there is no massless mode, and then there is no natural way to reduce the theory at the level of the fermionic field. That is, on the other hand, the case when $\phi = 0$, since it means that the $n = 0$ mode finds a natural $3 + 1$ dimensional interpretation and there is a smooth limit when $L \rightarrow 0$. Finally, parity breaking effects might be expected, only if there were a compelling reason to know that the gauge field itself adopts a topologically non-trivial configuration.

Acknowledgements

C.D.F. and C.C.T. thank CONICET for financial support. A.P.C.M. and I.R. thank CNPq/MCT and FAPERJ for partial financial support.

References

- [1] E. Brezin and J. Zinn-Justin, Nucl. Phys. B **257**, 867 (1985).
- [2] T. Kaluza, S. Preuss. Akad. Wiss. Berlin. (Math. Phys.), 966 (1921)
O. Klein, Z. Phys. Hadrons and Nuclei 37 (12): 895 (1926).
- [3] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **429**, 263 (1998)
- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999)

- [5] I. Antoniadis, Phys. Lett. B **246**, 377 (1990)
 I. Antoniadis, C. Muñoz, M. Quirós, Nucl. Phys. B **397**, 515 (1993)
 I. Antoniadis, S. Dimopoulos, A. Pomarol, M. Quirós, Nucl. Phys. B **544**, 503 (1999)
 I. Antoniadis, A. Pomarol, M. Quirós, Phys. Rev. D **60**, 095008 (1999)
- [6] K. R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. B **436**, 55 (1998)
- [7] L. J. Hall and Y. Nomura, Phys. Rev. D **65**, 125012 (2001)
- [8] M. Cvetič, G. Shiu and A.M. Uranga , Nucl. Phys. B **615**, 3 (2001)
- [9] T. Matsubara, Prog. Theor. Phys. 14 (1955) 351.
- [10] H. Ezawa, Y. Tomonaga and H. Umezawa, N. Cimento Ser. X, 5 810 (1957).
- [11] N.D. Birrell and L.H. Ford, Phys. Rev. D **22**, 330 (1980).
- [12] A. P. C. Malbouisson, J. M. C. Malbouisson and A. E. Santana, Nucl. Phys. B **631**, 83 (2002).
- [13] G. Panico and M. Serone, JHEP05, 024 (2005).
- [14] Y. Hosotani, Phys.Lett B **126**, 309 (1983); Phys.Lett B **129**, 193 (1983); Annals Phys. **190** 233 (1989).
- [15] See, for example, J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, Oxford Science Publications, 4th. Ed., (2002).
- [16] C. Fosco, G. L. Rossini and F. A. Schaposnik, Phys. Rev. Lett. **79**, 1980 (1997) [Erratum-ibid. **79**, 4296 (1997)] [arXiv:hep-th/9705124].